

RESEARCH

Open Access



Characterizing the neuroimmune environment of offspring in a novel model of maternal allergic asthma and particulate matter exposure

Juan M. Tamayo¹, Hadley C. Osman¹, Jared J. Schwartzer², Kent E. Pinkerton³ and Paul Ashwood^{1*}

Abstract

Inflammation during pregnancy is associated with an increased risk for neurodevelopmental disorders (NDD). Increased gestational inflammation can be a result of an immune condition/disease, exposure to infection, and/or environmental factors. Epidemiology studies suggest that cases of NDD are on the rise. Similarly, rates of asthma are increasing, and the presence of maternal asthma during pregnancy increases the likelihood of a child being later diagnosed with NDD such as autism spectrum disorders (ASD). Particulate matter (PM), via air pollution, is an environmental factor known to worsen the symptoms of asthma, but also, PM has been associated with increased risk of neuropsychiatric disorders. Despite the links between asthma and PM with neuropsychiatric disorders, there is a lack of laboratory models investigating combined prenatal exposure to asthma and PM on offspring neurodevelopment. Thus, we developed a novel mouse model that combines exposure to maternal allergic asthma (MAA) and ultrafine iron-soot (UIS), a common component of PM. In the current study, female BALB/c mice were sensitized for allergic asthma with ovalbumin (OVA) prior to pregnancy. Following mating and beginning on gestational day 2 (GD2), dams were exposed to either aerosolized OVA to induce allergic asthma or phosphate buffered saline (PBS) for 1 h. Following the 1-h exposure, pregnant females were then exposed to UIS with a size distribution of 55 to 169 nm at an average concentration of $176 \pm 45 \mu\text{g}/\text{m}^3$ (SD), or clean air for 4 h, over 8 exposure sessions. Offspring brains were collected at postnatal days (P)15 and (P)35. Cortices and hippocampal regions were then isolated and assessed for changes in cytokines using a Luminex bead-based multiplex assay. Analyses identified changes in many cytokines across treatment groups at both timepoints in the cortex, including interleukin-1 beta (IL-1 β), and IL-17, which remained elevated from P15 to P35 in all treatment conditions compared to controls. There was a suppressive effect of the combined MAA plus UIS on the anti-inflammatory cytokine IL-10. Potentially shifting the cytokine balance towards more neuroinflammation. In the hippocampus at P15, elevations in cytokines were also identified across the treatment groups, namely IL-7. The combination of MAA and UIS exposure (MAA–UIS) during pregnancy resulted in an increase in microglia density in the hippocampus of offspring, as identified by IBA-1 staining. Together, these data indicate that exposure to MAA, UIS, and MAA–UIS result in changes in the neuroimmune environment of offspring that persist into adulthood.

*Correspondence:

Paul Ashwood

pashwood@ucdavis.edu

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Keywords Neurodevelopment, Cytokines, Autism spectrum disorder (ASD), Schizophrenia, Neuroinflammation, Asthma/allergy, Fetal brain, Maternal immune activation (MIA), Maternal asthma and allergy (MAA)

Background

Neurodevelopmental disorders (NDD) are a broad group of disorders where central nervous system (CNS) development and function is altered [27]. NDD, including autism spectrum disorder (ASD) and attention deficit hyperactivity disorder (ADHD), are increasing in prevalence, with 1 in 6 children in the US experiencing a NDD in 2016 [125]. While there are some genetic origins of NDD, environmental factors such as exposure to pollution and maternal health are also recognized as risk factors [46, 100]. Mounting clinical and epidemiological data underscore the notion that environmental factors play a large role in both the underlying development of NDD and their impact on the severity of individual behavioral characteristics [52, 94, 99, 105, 114, 118, 120]. Pregnancy, in particular, marks a critical period when environmental insults have long-lasting effects on neurodevelopment.

The maternal immune system represents a key biological mechanism that links environmental factors to specific neurodevelopmental changes and later NDD diagnosis. For example, clinical studies have linked hospitalizations for bacterial and viral infections during pregnancy to an increased rate of birthing a child that will later be diagnosed with ASD [4, 14, 80, 116]. Moreover, laboratories have developed animal models that often use bacterial or viral mimics, such as lipopolysaccharide (LPS) or polyinosinic:polycytidylic acid (Poly(I:C)), respectively, to induce maternal inflammation during pregnancy and demonstrate a causal change in offspring brain and behavioral health [46]. Despite the efficacy of these models in demonstrating a link between maternal immune signaling and offspring neurodevelopment, they are limited in representing many of the common environmental sources of immune activation. With perhaps the exception of SARS-CoV-2, hospitalizations for viral infections are less frequent, and a recent meta-analysis of viral infections during pregnancy as a risk for children later being diagnosed with ASD was not supported [55]. Importantly, infections via bacterial and viral pathogens are only two of many ways in which the immune system can become activated, and researchers have begun to investigate other common environmental influences or immune conditions that can have a substantial impact on the maternal immune system and offspring brain development, notably allergic asthma.

Rates of asthma, like NDD, are currently increasing and represent a highly prevalent chronic disease that more commonly impacts ethnic/racial minority

and socioeconomically disadvantaged groups in the US [23, 72]. Importantly, chances of asthma exacerbations increase during pregnancy, and the presence of asthma increases likelihood of birthing a child that is later diagnosed with ASD [1, 3, 25, 26, 33, 42, 47, 87, 88, 93]. Asthma is an inflammatory disease of the lungs characterized by bronchial hyperresponsiveness, persistent inflammation, airflow obstruction, and reduction of airflow [39, 96]. The allergens associated with asthma can vary and include, but are not limited to rodent allergen, cockroach allergen, pollen, and house dust mite [53, 72]. Given the incidence of allergies and asthma in the United States [10, 34] and the high prevalence of allergic triggers in urban environments [22, 64, 72, 106], there is pressing need to better understand the causal effects of asthma during pregnancy and the impact of specific allergic triggers on offspring neurodevelopment.

Using our mouse model of maternal allergic asthma (MAA) to initiate an immune response in pregnant mice, we have previously reported systemic elevations in interleukin-4 (IL-4) and IL-5 in dams [109, 115] that parallel clinical reports associating increased IL-4 and IL-5 in mid-pregnancy maternal serum samples of mothers with children later diagnosed with ASD [40]. Not only do the dams in our MAA model produce a similar allergic asthma cytokine profile to that observed in humans, but the offspring display characteristic ASD-like behaviors, such as decreased social interaction and increased repetitive-like behaviors [20, 109, 110]. Moreover, MAA produces transcriptional differences in microglia gene expression and neuroinflammation in both prenatal offspring and brain regions of adult mice [20, 115, 117]. These findings highlight the lasting changes that allergic asthma during pregnancy can have on offspring neurodevelopment. However, environmental insults associated with ASD can vary, and exposure to these environmental factors does not necessarily occur in isolation for humans. In fact, individuals often simultaneously encounter multiple inflammatory-inducing environmental stimulants, and less is known about the potential synergistic effects of these exposures on maternal immune response and subsequent offspring development [22, 64, 106].

Particulate matter (PM), via air pollution, is another environmental factor that is suspected to be associated with the risk of neuropsychiatric disorders/NDD such as schizophrenia, attention deficit hyperactive disorder

(ADHD), and ASD [118–120]. Not only is PM linked to neurodevelopmental disruptions when exposures occur during pregnancy [21], but it has also been tied to exacerbated asthma responses and could potentially cause new onset cases of asthma [16, 17, 44, 108] although this remains controversial. PM is a complex mixture of constituents that contains organic compounds, soot, metals and metal-oxides, nitrates, and other elements in varying quantities depending on its source [113]. As such, it is necessary to characterize individual components of PM in order to effectively regulate air quality for limiting environmental exposure to toxicants.

Diesel exhaust is one common source of PM that has been identified as a risk factor for ASD [57, 103, 118–121]. Combustion-derived diesel exhaust includes soot particles containing elemental carbon and iron, which is the most common transition metal found in PM [127]. Iron in PM can occur from fuel additives and as a product of normal engine wear [74, 113], and iron-soot (IS) exposure has previously been shown to cause oxidative lung injury, induce inflammation of the lungs, and can cross the blood–brain barrier through nasal inhalation [48, 126, 127]. The links between PM and worsening of asthma symptoms and increased risk of neuropsychiatric disorders in offspring make PM exposure during pregnancy an important area that merits investigation.

There are many human studies and animal models investigating the links between gestational exposure to PM and neurodevelopmental outcomes in offspring [1, 2, 11, 118–120]. In addition, many studies have investigated the impacts of asthma as a risk factor for ASD, including our own [3, 20, 25, 26, 33, 42, 47, 68, 87, 88, 93, 109, 110, 115, 117]. However, these investigations consider asthma or PM only as independent risk factors for developing neuropsychiatric disorders. Despite the links between allergic asthma with ASD and the link between PM and ASD, there are no studies investigating the neuroimmune outcome on offspring of these environmental factors in conjunction. This is an apparent oversight given that PM exposure can worsen symptoms of asthma [44], and the source of environmental factors contributing to ASD are likely multifaceted.

In this study, we hypothesized that when mice are exposed to MAA and ultrafine iron-soot (UIS) particles during pregnancy (MAA–UIS), the neuroimmune outcomes will show heightened inflammation in offspring compared to that of MAA or UIS exposure alone. We also suspected that, because microglia can respond to changes in the neuroimmune environment through proliferation and are suspected to be associated with ASD behaviors [28, 51], we would see signs of functional differences in the frontal cortex and hippocampus—two brain regions implicated as being impacted developmentally in

neuropsychiatric disorders such as ASD [6, 31, 51, 102]. Using our established mouse paradigm of maternal aerosol exposure to study offspring outcomes, we demonstrate that MAA and UIS alone, as well as MAA and UIS combined, alter the neuroimmune profile in the brains of offspring that is sustained from adolescence into early adulthood.

Methods

Animals

BALB/c male and female mice were obtained from breeding pairs originally purchased from Envigo Laboratories (Livermore, CA, USA) and maintained at University of California, Davis at the Center for Health and Environment, Davis, CA. Mice were housed with same-sex littermates and kept at ambient room temperature (23 °C) on a 12 h light/dark cycle (lights on at 0800 h) within ventilated cages with water and food provided ad libitum. All procedures were performed with approval by University of California Davis Institutional Animal Care and Use Committee and according to guidelines established by National Institute of Health Guide for the Care and Use of Laboratory Animals.

Maternal allergic asthma and ultrafine soot particle exposure

Female BALB/c ($n=8$ /group) mice were sensitized with 10 μg of ovalbumin (OVA, Sigma, St. Louis, MO, USA) and 1 mg Al OH₃ dissolved in 200 μl of phosphate buffered saline (PBS) injected intraperitoneally at 7 and 8 weeks of age, for a total of two injections pre-pregnancy. Control dams were injected with PBS alone. Dams were then mated with age-matched males the week after final sensitization, and checked daily for the presence of a seminal plug, which was noted as gestational day (GD) 0.5. Beginning on gestational day 2 (G2), dams were exposed to either aerosolized OVA to induce allergic asthma or phosphate buffered saline (PBS) for 1 h, depending on treatment group. Following the 1-h exposure, mice were placed in a 20 cm \times 43 cm \times 18 cm polycarbonate whole-body chamber for exposure to an aerosol of ultrafine iron-soot (target concentration of 200 $\mu\text{g}/\text{m}^3$), including 40 $\mu\text{g}/\text{m}^3$ of iron oxide nanoparticles, or sham control clean air (AIR). The total iron-soot generated was cooled and diluted with filtered air to achieve the desired concentration prior to reaching the exposure chambers. Mice were exposed for 4 h/day on G2, G4, G6, G9, G11, G13, G16, and G18 to PBS or OVA [MAA condition] and AIR or ultrafine iron-soot [UIS condition], resulting in a total of four treatment groups: PBS/AIR (PBS-AIR), MAA/AIR (MAA), PBS/UIS (UIS), MAA/UIS (MAA–UIS). The mass concentration of the soot concentration was found by weighing

25-mm Teflon-coated filters (Teflo, Pall, East Hills, NY) on a microbalance before and after sample collection. The average soot concentration over the exposure period was $176 \pm 45 \mu\text{g}/\text{m}^3$ (SD). Following the last day of aerosol exposure (G18) pregnant mice were left undisturbed through parturition. Offspring were either sacrificed at postnatal day (P)15 or weaned at P21, housed with same-sex littermates, and sacrificed at P35 for brain tissue analysis.

Generation and characterization of particles

UIS particles were generated as previously described by Hopkins et al. [48]. Briefly, particle generation was obtained using a laminar diffusion flame system by mixing ethylene gas (the primary hydrocarbon fuel) and acetylene gas to compensate for the effect of iron oxide to suppress soot formation. By reaching the vapor phase of iron pentacarbonyl by warming to 20 °C with combusted argon carrier gas (all Sigma-Aldrich Chemical Co., St. Louis, MO) in the presence of an ethylene/acetylene vapor mix, the source of iron was generated. The result of these combusted reactants generated a hetero-disperse aerosol of ultrafine iron oxide particles (Fe_2O_3) and associated soot. Further details of the system and particle generation can be found in previously published studies [48, 54, 95]. In order to simulate unhealthy air quality conditions, an average particle concentration of 200 $\mu\text{g}/\text{m}^3$ was selected, as this reflects a concentration that is reminiscent of heavy pollution and poor air quality days in many parts of the world [48, 97].

Cardiac perfusion and brain tissue collections

At P15 and P35, one male and one female offspring from each litter were collected from their home cages, anesthetized using isoflurane (2–4% inhalation) and underwent transcardiac perfusion. Briefly, a lateral incision was made in the abdominal wall below the rib cage. With curved scissors, an incision was made in the diaphragm and cuts were made along the ribs to the collarbone to allow the sternum to be lifted. Once exposed, a 15-gauge perfusion needle was inserted into the ascending aorta of the heart for entry of perfusate, and an incision was made into the right atrium to create an outlet for drainage. Using a perfusion pump, 20 mL of PBS was slowly pumped through the circulatory system to reach adequate clearing. Whole brains were removed and dissected into hemispheric halves. One half was further dissected into cortical and hippocampal regions, flash frozen with liquid nitrogen, and stored at -80°C for later use in cytokine analyses. The remaining half was placed in 4% PFA for fixation for 24 h. Following this, it was then placed in 30% sucrose for 24 h for cryoprotection.

Cryoprotected tissues were then embedded in optimal cutting temperature (O.C.T.) media and frozen at -80°C .

Tissue sectioning and staining

Frozen tissue embedded in O.C.T. was sectioned with a Leica Instruments cryostat at 20 μm . Free-floating tissue sections were stored in PBS containing 0.01% sodium azide. Sections were incubated in 1:1000 rabbit-anti IBA-1 (Wako, Neuss, Germany) with 10% normal goat serum (NGS) and 0.2% triton X-100 at 4 °C for 24 h, followed by 1-h incubation with goat-anti rabbit biotinylated secondary antibody (Vector Laboratories, Burlingame, CA) in 5% NGS for 1 h at room temperature. Tissue sections were then incubated with avidin-biotinylated HRP complex (Vectastain Elite ABC kit, Vector Laboratories, Burlingame, CA) at room temperature. Visualization of labeling was conducted using 3,3'-diaminobenzidine (DAB) solution in the presence of peroxidase (HRP) enzyme. All sections were thoroughly rinsed three times with 1X PBS between staining steps. Sections were mounted onto glass Superfrost Plus microscope slides and cover slipped with VectaMount Permanent Mounting Medium.

Stereology

IBA-1-positive microglia were identified using stereological methods. IBA-1 cell counts were made on a bright-field microscope (Nikon Eclipse Ci, Nikon, Tokyo) at 20X magnification, and images were taken using NIS Elements v.4.0 (Nikon Instruments Inc. 1300 Walt Whitman Road Melville, NY 11747-3064, U.S.A.). Image analysis was performed using a macros script in ImageJ version 1.53 [98] to remove background noise and isolate IBA-1-positive cells in order to quantify microglia. A total of six to nine sections per brain region per animal were collected. Counts of microglia cells were taken from a 554.7 μm by 1232.1 μm box spanning the infralimbic and anterior cingulate cortical areas of the frontal cortex, as well as the dentate gyrus, CA1, CA2, and CA3 of the hippocampus. Microglia were identified by IBA-1 positive cell body staining. Statistical significance was determined using one-way ANOVA with multiple comparisons on GraphPad Prism version 9.4.1 (GraphPad software, graphpad.com).

Multiplex bead-based cytokine analysis

Frozen tissues were lysed using cell lysis buffer and total protein from the lysates were quantified via Bradford assay. Analysis of serum cytokines was performed using a multiplex mouse 25-plex bead immunoassay (Milliplex Mouse Cytokine/Chemokine Magnetic Bead Panel #MCYTMAG70PMX25BK). The following

cytokines were quantified: granulocyte colony stimulating factor (G-CSF), granulocyte colony stimulating factor (GM-CSF), IFN- γ , IL-1 α , IL-1 β , IL-2, IL-4, IL-5, IL-6, IL-7, IL-9, IL-10, IL-12 (p40), IL-12 (p70), IL-13, IL-15, IL-17, interferon gamma-induced protein 10 (IP-10), keratinocyte chemoattractant (KC), monocyte chemoattractant protein-1 (MCP-1), macrophage inflammatory protein-1 alpha (MIP-1 α), MIP-1 β , MIP-2, chemokine ligand 5 (CCL5/RANTES), and tumor necrosis factor alpha (TNF- α). Standards and reagents were all prepared according to the manufacturers' recommendations. Each lysate was diluted to a standardized concentration of 50 μ g and run in duplicate. Twenty-five microliters of sample, standards, or blanks were loaded into a 96-well plate with appropriate amounts of assay buffer and matrix solution. The plate was then incubated overnight with antibody-coupled magnetic beads. The following day, the plate underwent a series of washes. Washes were performed using a Bio-Plex handheld magnet (Bio-Rad Laboratories, Hercules, CA, USA). After the final wash, the plate was incubated with biotinylated detection antibody on a shaker for 30 min, and analyzed using a Bio-Rad Bio-Plex 200 plate reader (Bio-Rad Laboratories, Hercules, CA, USA). The following were the minimal amounts of detectable cytokine concentration: G-CSF: 1.7 pg/mL; GM-CSF: 10.9 pg/mL; IFN γ : 1.1 pg/mL; IL-1 α : 10.3 pg/mL; IL-1 β : 5.4 pg/mL; IL-2: 1.0 pg/mL; IL-4: 0.4 pg/mL; IL-5: 1.0 pg/mL; IL-6: 1.1 pg/mL; IL-7: 1.4 pg/mL; IL-9: 17.3 pg/mL; IL-10: 2.0 pg/mL; IL-12 (p40): 3.9 pg/mL; IL-12 (p70): 4.8 pg/mL; IL-13: 7.8 pg/mL; IL-15: 7.4 pg/mL; IL-17: 0.5 pg/mL; IP-10: 0.8 pg/mL; KC: 2.3 pg/mL; MCP-1: 6.7 pg/mL; MIP-1 α : 7.7 pg/mL; MIP-1 β : 11.9 pg/mL; MIP-2: 30.6 pg/mL; RANTES: 2.7 pg/mL; TNF- α : 2.3 pg/mL. Sample concentrations that fell below the minimal detection value were given a proxy value of half the limit of detection for statistical comparisons.

Statistical analysis

Brain cytokine data were analyzed using linear-mixed effects modeling to control for the unexplained residual variance that could be originating from litter-to-litter variations due to the hierarchical data structure in which statistical independence of observation is violated [12, 61, 62, 107, 122]. Models were built with the lme package of R version 4.2.2 using a forward-stepwise regression approach. First, a random-effects only model was constructed with litter set as the random effect. Then, two fixed effects variables were added for the exposure types: MAA (OVA) and UIS (Soot). A third model was constructed that included both the main effects and the interaction of MAA and UIS followed by a final model containing both treatments, sex, and their interactions. Model fit was assessed using the likelihood ratios test

and the best model was selected based on the Akaike Information Criterion (AIC). For models with significant interactions effects, groups were further analyzed using pairwise comparisons of estimated marginal means. Litter size and ratio of male offspring were assessed by two-way analysis of variance (ANOVA).

Results

Litter size and sex ratio

Litter sizes averaged 5–6 offspring per litter (see Table 1), with no differences in the number of pups for MAA ($p=0.757$), UIS ($p=0.925$), or their interaction ($p=0.683$). Similarly, no differences were observed between the male-to-female ratio across treatment groups ($p > 0.05$ for all comparisons).

P15 offspring cortex cytokines

Offspring brains were collected and micro-dissected into cortical and hippocampal sections at P15. Homogenates of each section were then analyzed for cytokine concentration. Multilevel mixed-effects modeling was used to control for within-litter variability and inclusions of fixed effects were determined using forward stepwise regression. For all cytokines measured, the inclusion of sex-difference did not significantly improve model fit (Additional file 1: Table S1). In the cortex, several cytokines were elevated in both male and female offspring of MAA, UIS, and MAA–UIS dams, many of which are generally considered inflammatory and potentially neurotoxic with prolonged exposure (Fig. 1). Among these, three cytokines (IL-1 β , IL-2, IL-17) were found to be significantly elevated in all treatment groups. For example, a main effect of exposure for both MAA ($p=0.016$), and UIS ($p < 0.001$), and MAA–UIS ($p < 0.001$) resulted in an average of a 2- to threefold increase of IL-1 β in the cortex, compared to age-matched PBS-AIR controls (Fig. 1). No additional additive effects were observed with the combination of MAA and UIS. Similarly, it was found that there were significant increases in IL-2 for both MAA ($p=0.002$) and UIS ($p=0.001$) exposures. For IL-2, the MAA–UIS interaction model identified a significant elevation of IL-2 in the MAA–UIS

Table 1 Litter size and sex ratio

Treatment	Litter size	Ratio of males
PBS/air	5.83 \pm 1.83	0.49 \pm 0.15
MAA	5.40 \pm 1.14	0.34 \pm 0.20
UIS	5.50 \pm 1.29	0.33 \pm 0.13
MAA–UIS	5.60 \pm 1.14	0.49 \pm 0.23

Ratio of male-to-female offspring was calculated as the number of male mice in a litter divided by the total number of pups in the litter. Litter size and sex ratio data are shown as mean \pm SD




Fig. 1 Cortical cytokine concentrations in P15 offspring brains exposed to PBS-AIR, MAA, UIS, or MAAUIS. Cytokines were assessed using multiplex bead-based immunoassays. **A** IL-1 β , **B** IL-2, **C** IL-17, **D** IL-13, **E** MIP-1 α , **F** IP-10, **G** IL-7 are represented as pg/mL after being normalized to total protein content. Statistical significance was determined by multilevel mixed-effects modeling. Main effects are represented as * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ compared to PBS-Air; pairwise comparisons are represented as # $p < 0.05$, ## $p < 0.01$

group compared to the PBS-Air group ($p=0.045$). However, post hoc analysis did not identify any interaction effects between MAA–UIS and the other treatment conditions. Similarly, a model containing the MAA–UIS interaction for IL-17 confirmed main effects for both MAA ($p=0.001$) and UIS ($p<0.001$) exposures, as well as a significant MAA–UIS interaction ($p=0.017$), but the differences between these treatment conditions were not statistically significant.

The elevation in IL-1 β , IL-2, and IL-17 occurred with concomitant increases in the canonical anti-inflammatory marker IL-10. A model containing treatment conditions and their interaction identified effects of both MAA ($p=0.001$), and UIS ($p=0.005$) on IL-10 elevations compared to PBS-Air controls. Interestingly, a significant MAA–UIS interaction ($p=0.002$) identified suppressive effects of the combined MAA–UIS exposure on IL-10 levels. Post hoc analysis confirmed that while the MAA ($p=0.007$) and UIS ($p=0.024$) alone both increased the IL-10 concentration compared to PBS-Air, this increase was absent in the MAA–UIS group ($p=0.387$) and no differences were observed between MAA and UIS conditions ($p=0.967$).

For several inflammatory markers, only the main effect of UIS significantly altered cytokine concentrations in the cortex. Specifically, the exposure to UIS resulted in an approximately 54 pg/mL ($p<0.001$), and 49 pg/mL ($p=0.003$) increase in IL-13 and MIP-1 α , respectively, compared to the PBS-Air group, and no effect of MAA were observed for either of these two cytokines. Conversely, a main effect of MAA was observed for IL-7 ($p=0.050$) without an effect of UIS ($p=0.167$). For both IL-13, MIP-1 α , and IL-7, models containing a MAA–UIS interaction were not significant and no sex-by-treatment interactions were identified in any of the P15 cortex cytokine measures.

P35 offspring cortex cytokines

Littermates from the P15 cytokine investigation were left undisturbed until P35, at which time brains were collected and micro-dissected into cortical and hippocampal sections. Mixed-effects models that included sex were not significantly improved over treatment-alone models for all cytokines measured (Additional file 2: Table S2). However, models that included treatment main effect and interaction revealed changes in several cytokines in the cortices of both sexes of MAA, UIS, and MAA–UIS offspring at P35 in a similar manner to what was identified in P15 offspring (Fig. 2). Many of the cytokine changes observed at P15 remained altered in the P35 offspring cortex. For instance, it was revealed that there were effects of MAA and UIS in IL-1 β , IL-13 and IL-17 cortex of P35 offspring compared to PBS-AIR control

mice (IL-1 β : MAA $p=0.046$, UIS $p=0.008$; IL-13: MAA $p=0.002$, UIS $p=0.005$; IL-17: MAA $p<0.001$, UIS $p=0.001$) (see Additional file 2: Table S2). While mixed-effects models were selected that included an MAA–UIS interaction for these cytokines, post hoc analysis revealed no additive or suppressive effects of the combined exposure of MAA and UIS. Elevations in IL-1 β , IL-13, and IL-17 were similar between both MAA- and UIS-alone groups ($p>0.05$ for all comparison). Similarly, main effects for both MAA and UIS exposure were observed in the P35 cortex for KC, IL- α , and MIP-1 α (KC: MAA $p=0.002$, UIS $p=0.050$; IL-1 α : MAA $p=0.050$, UIS $p=0.013$; MIP-1 α : MAA $p=0.002$, UIS $p=0.021$). No differences were observed in any of these cytokine concentrations between in the three exposure groups and no interactions for MAA–UIS were observed in the mixed effects models.

For several cytokines the individual contribution of MAA, but not UIS, impacted specific inflammatory profiles. Specifically, MAA treatment resulted in an increase in IL-9 ($p=0.012$), IL-4 ($p=0.037$), and IL-2 ($p=0.005$) compared to offspring born from PBS-Air exposed dams. Of all the cytokines investigated, IP-10 concentration was the only identified cytokine where a combination of MAA and UIS resulted in suppressive effects. Specifically, the mixed-effects model revealed that MAA exposure alone significantly increased the level of IP-10 in the P35 cortex compared to PBS-Air offspring, with an additional interaction identified in the combination of MAA and UIS for IP-10 ($p=0.008$) suggesting a suppressive effect by UIS. Post hoc analysis confirmed that MAA-only offspring had significantly higher levels of IP-10 compared to MAA–UIS mice ($p=0.026$), and no differences were observed between the MAA–UIS condition and PBS-Air control mice ($p=0.977$).

P15 offspring hippocampus cytokines

Along with cortical sections, hippocampal homogenates from the same offspring brains were analyzed at P15 for cytokine concentrations. For all cytokines measured, the inclusion of sex in multi-level models did not significantly improve model fit (Additional file 3: Table S3). Similar to what was uncovered in the respective cortical sections, several inflammatory cytokines were elevated in male and female offspring of MAA, UIS, and MAA–UIS dams (Fig. 3). Interestingly, main effects for both MAA ($p=0.002$) and UIS ($p=0.005$) were only observed for IL-7 in the P15 hippocampus. None of the other cytokines investigated reached statistical significance in all three treatment groups compared to controls. However, IFN γ concentrations showed a main effect for UIS ($p=0.007$) but not MAA ($p=0.111$), and similar elevations were observed in IL-12(p40) and IL-17 for UIS




Fig. 2 Cortical cytokine concentrations in P35 offspring brains exposed to PBS-Air, MAA, UIS, or MAAUIS. Cytokines were assessed using multiplex bead-based immunoassays. **A** IL-1 β , **B** IL-13, **C** IL-17, **D** KC, **E** IL-1 α , **F** MIP-1 α , **G** IL-9, **H** IL-4, **I** IL-2, and **J** IP-10 are represented as pg/mL after being normalized to total protein content. Statistical significance was determined by multilevel mixed-effects modeling. Main effects are represented as * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ compared to PBS-Air




Fig. 3 Hippocampal cytokine concentrations in P15 offspring brains exposed to PBS-AIR, MAA, UIS, or MAAUIS. Cytokines were assessed using multiplex bead-based immunoassays. **A** IL-7, **B** IFN γ , **C** IL-12(p40), **D** IL-17, **E** IL-13, **F** IL-1 β , **G** IL-2, **H** RANTES, and **I** IL-15 are represented as pg/mL after being normalized to total protein content. Statistical significance was determined by multilevel mixed-effects modeling. Main effects are represented as * $p < 0.05$, ** $p < 0.01$, compared to PBS-Air; pairwise comparisons are represented as # $p < 0.05$, ## $p < 0.01$

alone or with MAA–UIS (IL-12(p40): $p < 0.007$; IL-17: $p < 0.002$). These increases were not present in the hippocampus of offspring born from the MAA treatment-alone group. Interestingly, many cytokines measured in the P15 hippocampus showed suppressive effects of the combined treatment of MAA and UIS. For example, a mixed-effects model for IL-13 containing a treatment interaction revealed a significant effect of MAA-alone ($p = 0.034$) and UIS-alone ($p = 0.011$) as well as a negative effect of the MAA–UIS interaction ($p = 0.014$). Post hoc comparisons confirmed that the IL-13 concentrations in the MAA–UIS offspring hippocampus were no different from PBS-Air mice ($p = 0.272$). However, differences between MAA-alone or UIS-alone groups were also not statistically different from the MAA–UIS condition ($p > 0.05$ for all comparisons). Similarly, these intermediate effects were also observed for IL-1 β (MAA $p = 0.047$, UIS $p = 0.038$), IL-2 (MAA $p = 0.026$, UIS $p = 0.002$), and RANTES (MAA $p = 0.041$, UIS $p = 0.034$). Finally, one additional cytokine was found to be impacted by MAA alone compared to the control, and not the UIS or MAA–UIS treatment groups. Specifically, IL-15 was found to be elevated by MAA ($p = 0.009$), but not UIS or MAA–UIS.

P35 offspring hippocampus cytokines

Hippocampal homogenates from P35 offspring were also assessed for cytokine differences; however, multi-level mixed-effects modeling did not reveal any significant effects of treatment or sex in on the concentration of all cytokines measured ($p > 0.05$; see Additional files 1, 2, 3, 4).

P15 Microglia density

Microglia within a 554.7 μm by 1232.1 μm box spanning the anterior cingulate cortex, prelimbic area, and infralimbic area were counted using ImageJ by an investigator blinded to treatment condition. Analysis of microglia density in the dentate gyrus, CA1, CA2, and CA3 regions of the hippocampus in P15 mice showed significant differences between the PBS-air group and the MAA–UIS group ($p = 0.0166$; Fig. 4). There were no significant changes in the microglia density between treatment groups in the frontal cortex of P15 mice (Fig. 4). Additionally, in both the hippocampus and cortex, no differences in microglia density were seen in the single treatment groups (MAA or UIS) when compared to the PBS-air group.

Discussion

Among the many well established environmental factors that can impact fetal neurodevelopment, asthma and air pollution represent two major sources of immune stimulation that are on the rise, making them a significant

concern for pregnant individuals. Based on previous studies of maternal immune activation with asthma, and PM exposure during pregnancy, we hypothesized that the combination of these two environmental stimuli would result in an exacerbated neuroimmune response in offspring. Although the appearance of an additive or synergistic effect of MAA and UIS exposure combined was limited, we did identify increases in cytokine concentrations across all treatment groups in the cortex and hippocampus that may suggest converging pathways for each insult/exposure. However, we also observed potentially suppressive activities that may suggest activation of competing or diverging pathways. Importantly, some of changes appear to be sustained across treatment groups from adolescence into early adulthood in the cortex, demonstrating lasting impacts of these gestational exposures on the neuroimmune environment later in life. Although we identified increases in cytokines in the hippocampus within all treatment groups at P15, these did not remain elevated into early adulthood. Overall, these data show that MAA and UIS environmental stimuli can result in an altered neuroimmune environment that persists from juvenile to adult timepoints.

The allergic response in the lungs of individuals with asthma is characterized by an influx of immune cells, such as neutrophils, mast cells, macrophages, and T-helper (T_H)2 cells. Our mouse model of MAA previously showed elevated IL-4, IL-5, IL-17, and IFN γ in the lung and peripheral blood of mice exposed to aerosolized OVA during pregnancy [20, 110, 115]. These increases in maternal serum cytokines result in neuro-immune signaling changes in the fetal brain during in utero development [115]. Our present data extend these findings by revealing evidence of increases in cortical and hippocampal cytokines in juvenile mice of MAA dams. In the cortex, for example, MAA increased IL-1 β , IL-2, IL-17, IL-10, and IL-7. In the hippocampus, we identified IL-1 β , IL-2, IL-7, IL-13, IL-15, and RANTES as being elevated in P15 offspring of MAA-exposed dams. These observed increases in MAA compared to PBS-AIR controls demonstrate the independent neurodevelopmental impact of allergic inflammation during pregnancy on offspring neuroinflammation. In addition to these findings in the MAA-only treatment group, we also identified suppressive effects of combining MAA with UIS in the cortex in juvenile offspring, notably, IL-10 an anti-inflammatory cytokine, whose suppression may suggest a lack of immune regulation and shift towards more neuroinflammation. Moreover, increases in cytokines as a result of UIS exposure alone were also identified in the hippocampus of juvenile offspring, with elevated IFN γ , IL-1 β , IL-7, and IL-12(p40). To the authors' knowledge, investigations into these neurobiological outcomes in offspring under




Fig. 4 Microglia density in the hippocampus and frontal cortex of MAA, UIS, and MAA-UIS of P15 offspring. **a** and **c** Staining for IBA-1 with diaminobenzidine (DAB) was used to label microglia in coronal sections of p15 mice in the **(a)** frontal cortex and **(c)** hippocampus. **b** and **d** Quantification of microglial density in the **(b)** frontal cortex and **(d)** hippocampus. Statistical significance determined via one-way ANOVA. In the frontal cortex, $n=6$ (PBS-AIR), $n=4$ (MAA), $n=5$ (UIS), $n=3$ (MAA-UIS). In the hippocampus, $n=9$ (PBS-AIR), $n=7$ (MAA), $n=6$ (UIS), $n=6$ (MAA-UIS)

UIS exposure during gestation have not been previously reported, highlighting the novelty of our model and findings.

In addition to independent effects of MAA or UIS treatment on cytokine concentrations in the cortex and hippocampus of juvenile offspring, several cytokines did not respond further to the combined effects of MAA and UIS. Most notably, we observed similar elevations in the

cortex of IL-1 α , IL-1 β , IL-2, and IL-17 in dual-exposed MAA-UIS offspring suggesting a potential ceiling effect produced by either environmental insult alone, or a maximal response through converging signaling pathways. IL-1 β , IL-13, and IL-17 were also trending or increased in the MAA and UIS single treatment groups as well as the MAA-UIS treatment group at P15, and they remained elevated into the P35 timepoint in the cortex. Our data

suggest a sustained elevation in these cytokines from P15 to P35 as a result of both MAA and UIS that have the potential for long-lasting impacts on neurodevelopment in the cortex of these offspring.

Consistent with the pleiotropic nature of cytokines in the central nervous system (CNS), IL-1 β , IL-2, and IL-17 have all been identified as having neurotrophic properties [5, 13, 29, 58, 66, 69, 78, 92] (Rochman et al. 2013). Indeed, high concentrations (500 ng/mL) of IL-1 β can have neurotoxic effects on neurons when exposed for 3–5 days [92], and IL-17 is detected at high levels in the CNS in multiple sclerosis and associated with the neuroinflammatory pathology of the disease [58, 66]. Compared to these neurotoxic concentrations, our data represent moderate increases in cytokines with less than 2.5-fold increases in treatment groups compared to PBS-AIR controls, and may not represent overt inflammation per se. However, these smaller changes in brain cytokine levels during the juvenile period may be biologically significant given their alternative functions in promoting neuronal survival and neurodevelopment. For example, IL-1 β acts as a chemokine that guides neurite outgrowth in cortical neurons [69], and IL-17 acts in initiating the release of brain-derived neurotrophic factor (BDNF), glia-derived neurotrophic factor (GDNF), and nerve growth factors (NGF) that are associated with neuronal cell survival and repair [78]. Taken from this view, these cytokines, which are generally considered overtly inflammatory, may be having a more subtle impact on the neuroarchitecture of offspring brains than models finding dramatic increases in concentration of IL-1 β and IL-17.

Further demonstrating the neuropoietic nature of these cytokines within the CNS and adding to the idea that the moderate increases observed in this model may be altering the neuropatterning of the offspring brains, IL-2 has been found to have neurotrophic properties and is necessary for proper cytoarchitecture in development [8, 8, 9, 9]. Both IL-2 and IL-13 are among several cytokines that are known to decrease in concentration at P14 under homeostatic conditions, and this developmental time-point in mice is characterized by a high degree of synaptogenesis and pruning [37, 81]. In contrast, our model, which investigated cytokine concentrations at P15, still within this window of high synaptogenesis, showed increased IL-2, representing a shift in homeostatic load. Taken together, it may be that these sustained moderate increases in cytokines of the cortex are changing the trajectory of cortical development and promoting altered connectivity in the cortex linked to behavioral changes such as decreased social interaction and repetitive behaviors previously identified in our model [19, 109]. This phenomenon of altered connectivity has also been implicated in the core behaviors associated with ASD,

specifically the social deficits and restrictive and repetitive behaviors [24, 73, 77].

Similar to our findings in the cortex, we also identified elevations in several cytokines, most notably IFN γ and IL-7, at P15 in the hippocampus of MAA and/or UIS groups compared to control offspring. IFN γ receptors are present on both neurons and glia [90]. In the hippocampus, IFN γ appears to play a role in synaptogenesis [65]. Some investigators have found that overexpression resulted in increased neurogenesis in the dentate gyrus, and because of its neuromodulatory effects, it has been suggested that this may impact cognition and social behavior as a result [7, 35, 36]. Additionally, IL-7 promotes survival and neurite outgrowth in hippocampal neurons [70, 76]. Considering the effects of IL-7 and IFN γ , and that the hippocampus is a major neurogenic niche in the developing brain, future studies may benefit from investigating the potential for hippocampal overgrowth in offspring brains in response to UIS or allergic asthma inflammation during pregnancy. Indeed, this phenomenon of hippocampal overgrowth has been identified in cases of ASD [43, 86, 104] and has been implicated in the deficits associated with emotion perception and sensory processing in ASD individuals [6, 43]. Curiously, the observed increases in hippocampal cytokine concentrations at P15 were not observed in any treatment group of P35 offspring compared to PBS-AIR controls. Although we can only speculate about these findings, it may be that these changes resolve during adolescence when additional brain maturation may be present to compensate for developmental overgrowth much in the same way that volumetric increases in the hippocampus of ASD individuals do not persist into adulthood [43, 104]. Although IL-7 and IFN γ are only two examples of cytokines that were elevated across the treatment groups, they illustrate the broader findings that treatment with MAA, UIS, or MAA–UIS can alter the hippocampal neuroimmune environment.

Prenatal insults have been shown to have lifelong impacts on microglia function and are suspected to play a prominent role in NDD [32, 38, 50, 59, 112, 117]. In ASD, some postmortem studies have identified differences in microglia density and morphology in brains of individuals [60, 82, 83]. In our previous study of MAA, we found DNA methylation differences in adult microglia, and several of these changes occurred in regulatory genes that are shared among some ASD individuals [117]. Given these findings, we sought to examine the density of microglia in the P15 brains of our MAA and UIS exposure model. We observed a significant increase in microglia density within the hippocampus of offspring exposed to MAA–UIS, but these increases were not present in the frontal cortex. One plausible explanation for

why these increases were only observed in the hippocampus may be due to the higher density of microglia known to be present in the hippocampus. This higher density of microglia is thought to make the hippocampus more vulnerable to inflammation [18, 101], and disruptions in the dentate gyrus have been linked to NDD [15, 124]. Our findings of increased microglia density in the hippocampus of MAA–UIS offspring mirror data from another maternal immune activation model that utilizes the viral mimic poly(I:C). Specifically, Juckel et al. [56] reported an increase in microglia density in the hippocampus but not the cortex of offspring born from immune-activated dams. Similarly, another study of maternal immune activation using LPS stimulation showed an increase in microglia density in the hippocampus [30]. While it is difficult to make conclusions about microglial function based on density data alone, our observed difference in the hippocampus in combination with similar reports from other maternal immune activation models [30, 56], suggest that asthma allergy and PM mediated immune activation during pregnancy can result in a deviation from homeostatic activity in the offspring hippocampus. Of interest, IFN γ and IL-12 were increased in MIA-UIS in the hippocampus, cytokines that are often associated with a T_H1 response and M1 macrophage/microglia skewing. These cytokines were not elevated in the cortex. Moreover, in the cortex and hippocampus UIS is associated with increased IL-13 that may balance the signals from IL-12 and IFN γ as seen in peripheral responses. IL-13 is often considered anti-inflammatory in the CNS, with some studies pointing to a neuroprotective impact in CNS diseases and injuries [45, 63, 75, 91]. Although speculative, the results may suggest combined inflammatory M1/T_H1 associated cytokines IFN γ and IL-12, could influence the MAA–UIS microglia density results while these are protected in the cortex or under UIS conditions through IL-13 production. More investigation is needed on the combination of cytokines released and action on microglia cells in the context of environmental exposures.

Although we did not collect maternal data in this preliminary study, data from previous MAA studies demonstrate increased systemic inflammation characteristic of an allergic asthma response, specifically with increased IL-4, IL-5, and IL-13 [19, 110, 115], suggesting the potential for a similar response in dams of MAA in the current model. Speculation about the systemic impacts of UIS on the maternal immune system, however, is difficult. Many studies of PM exposure suggest IL-6, IL-8, and TNF α as the main cytokines upregulated in response to PM exposure [41, 67, 79, 89, 111]. This difference in cytokine response highlights the potential reason we see differences in the impact between MAA and UIS in our model. However, models of PM can vary widely in the size of

PM and composition [79], making speculation about the maternal response in the UIS groups, and the potential role this plays in offspring neurodevelopment, difficult. This variation in PM studies underscores the need for future investigations to identify the maternal cytokine milieu in this model.

While our findings do not necessarily demonstrate an additive effect of MAA and UIS with regard to the cytokines we investigated, we did see a synergistic impact of MAA–UIS on microglia density in the hippocampus. These findings demonstrate the potential for additive effects of maternal asthma exposure when coupled with PM exposure. Independently, studies have shown in both humans and animal models that PM exposure during pregnancy can increase the susceptibility of offspring developing asthma [49, 84, 123]. This increased susceptibility of asthma in offspring is also seen in children of asthmatic mothers [71, 85] (Mattes et al. 2013), suggesting the potential for systemic immune dysfunction when these two stimuli are combined during pregnancy. The findings in this unique model of MAA and UIS exposure highlight the importance of investigating the impact of these closely linked and prevalent environmental factors.

Conclusions

Our data add to our previous studies on the impact of MAA on fetal brain development, showing here that this model impacts region-specific cytokine concentrations in both the juvenile and adolescent periods. To the investigators' knowledge, this was the first study to assess the impact of ultrafine iron-soot exposure during gestation on offspring neurobiology. Taken together, these data highlight the importance of understanding the impact that common environmental stimuli can have on fetal development, and the potential for these stimuli to have long-lasting changes in offspring.

Abbreviations

ASD	Autism spectrum disorders
PM	Particulate matter
MAA	Maternal allergic asthma
UIS	Ultrafine iron-soot
OVA	Ovalbumin
GD2	Gestational day 2
PBS	Phosphate buffered saline
P	Postnatal days
IL	Interleukin
IFN γ	Interferon gamma
MAA–UIS	MAA and UIS exposure
LPS	Lipopolysaccharide
Poly(I:C)	Polyinosinic:polycytidylic acid
ADHD	Attention deficit, hyperactive disorder
IS	Iron-soot
NGS	Normal goat serum
DAB	3,3'-Diaminobenzidine
G-CSF	Granulocyte colony stimulating factor
GM-CSF	Granulocyte colony stimulating factor

IP-10	Interferon gamma-induced protein 10
KC	Keratinocyte chemoattractant
MCP-1	Monocyte chemoattractant protein-1
MIP-1 α	Macrophage inflammatory protein-1 alpha
RANTES	Chemokine ligand 5
TNF- α	Tumor necrosis factor alpha
AIC	Akaike information criterion
T _H	T-helper
CNS	Central nervous system
BDNF	Brain-derived neurotrophic factor
GDNF	Glia-derived neurotrophic factor
NGF	Nerve growth factors
MIA	Maternal immune activation

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12974-023-02930-7>.

Additional file 1. Table S1.

Additional file 2. Table S2.

Additional file 3. Table S3.

Additional file 4. Table S4.

Acknowledgements

We would like to thank Mr. Dale Uyeminami and the undergraduate interns who contributed to this work.

Author contributions

JMT participated in overall project conceptualization, participated in and managed mouse colony husbandry, performed tissue preparation, experiment design for all experiments, data collection, analysis, interpretation for Luminex assay experiments and microglia density assay for hippocampal tissue, manuscript drafting and revisions. HCO acquired, analyzed, and interpreted data for cortical microglia density assay, drafted results for microglia density assay, contributed to revisions. JJS reviewed and edited manuscript and consulted on statistical analysis. KP participated in conceptualization, methodology, project administration, supervision, manuscript revisions and editing, funding acquisition. PA designed the study, methodology, project administration, supervision, contributions to interpretation of the data, manuscript revisions and editing, funding acquisition. All authors read and approved the final manuscript.

Funding

This material is based upon work supported by National Institute of Environmental Health Sciences (R21ES025560, R21ES035492, P30 ES023513 and NIOSH U54 OH07550), and the National Institute of Mental Health (R21MH116383, R01MH118209, R15MH119500).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The studies involved animal models and were reviewed and approved by University of California, Davis Institutional Animal Care and Use Committee (IACUC) and according to guidelines established by National Institute of Health Guide for the Care and Use of Laboratory Animals.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Medical Microbiology and Immunology, and the M.I.N.D. Institute, University of California at Davis, 2805, 50th Street Sacramento, Davis, CA 95817, USA. ²Program in Neuroscience and Behavior, Department of Psychology and Education, Mount Holyoke College, 50 College Street, South Hadley, MA 01075, USA. ³Center for Health and the Environment, University of California at Davis, Davis, CA 95616, USA.

Received: 4 July 2023 Accepted: 12 October 2023

Published online: 02 November 2023

References

1. Abdallah MW, Larsen N, Grove J, Nørgaard-Pedersen B, Thorsen P, Mortensen EL, Hougaard DM. Amniotic fluid inflammatory cytokines: potential markers of immunologic dysfunction in autism spectrum disorders. *World J Biol Psychiatry*. 2013;14(7):528–38.
2. Ahadullah, Yau SY, Lu HX, Lee TMC, Guo H, Chan CCH. PM2.5 as a potential risk factor for autism spectrum disorder: its possible link to neuroinflammation, oxidative stress and changes in gene expression. *Neurosci Biobehav Rev*. 2021;128: 534–548.
3. Ali Z, Ulrik CS. Incidence and risk factors for exacerbations of asthma during pregnancy. *J Asthma Allergy*. 2013;6:53–60.
4. Atladóttir HO, Thorsen P, Østergaard L, Schendel DE, Lemcke S, Abdallah M, Parner ET. Maternal infection requiring hospitalization during pregnancy and autism spectrum disorders. *J Autism Dev Disord*. 2010;40(12):1423–30.
5. Awatsuji H, Furukawa Y, Nakajima M, Furukawa S, Hayashi K. Interleukin-2 as a neurotrophic factor for supporting the survival of neurons cultured from various regions of fetal rat brain. *J Neurosci Res*. 1993;35(3):305–11.
6. Banker SM, Gu X, Schiller D, Foss-Feig JH. Hippocampal contributions to social and cognitive deficits in autism spectrum disorder. *Trends Neurosci*. 2021;44(10):793–807.
7. Baron R, Nemirowsky A, Harpaz I, Cohen H, Owens T, Monsonego A. IFN-gamma enhances neurogenesis in wild-type mice and in a mouse model of Alzheimer's disease. *FASEB J*. 2008;22(8):2843–52.
8. Beck RD Jr, King MA, Ha GK, Cushman JD, Huang Z, Petitto JM. IL-2 deficiency results in altered septal and hippocampal cytoarchitecture: relation to development and neurotrophins. *J Neuroimmunol*. 2005;160(1–2):146–53.
9. Beck RD Jr, Wasserfall C, Ha GK, Cushman JD, Huang Z, Atkinson MA, Petitto JM. Changes in hippocampal IL-15, related cytokines, and neurogenesis in IL-2 deficient mice. *Brain Res*. 2005;1041(2):223–30.
10. Blackwell DL, Lucas JW, Clarke TC. Summary health statistics for U.S. adults: national health interview survey, 2012. *Vital and health statistics. Series 10, Data from the National Health Survey*. 2014; (260), 1–161.
11. Bolton JL, Marinero S, Hassanzadeh T, Natesan D, Le D, Belliveau C, Mason SN, Auten RL, Bilbo SD. Gestational exposure to air pollution alters cortical volume, microglial morphology, and microglia-neuron interactions in a sex-specific manner. *Front Synaptic Neurosci*. 2017;9:10.
12. Brauer M, Curtin JJ. Linear mixed-effects models and the analysis of nonindependent data: a unified framework to analyze categorical and/or continuous independent variables that vary within-subjects and/or within-items. *Psychol Methods*. 2018;23(3):389–411.
13. Brombacher TM, Nono JK, De Gouveia KS, Makena N, Darby M, Womersley J, et al. IL-13-mediated regulation of learning and memory. *J Immunol*. 2017;198(7):2681–8.
14. Brown AS, Sourander A, Hinkka-Yli-Salomäki S, McKeague IW, Sundvall J, Surcel HM. Elevated maternal C-reactive protein and autism in a national birth cohort. *Mol Psychiatry*. 2014;19(2):259–64.
15. Cai Y, Tang X, Chen X, Li X, Wang Y, Bao X, et al. Liver X receptor β regulates the development of the dentate gyrus and autistic-like behavior in the mouse. *Proc Natl Acad Sci*. 2018;115(12):E2725–33.
16. Chalupa DC, Morrow PE, Oberdörster G, Utell MJ, Frampton MW. Ultrafine particle deposition in subjects with asthma. *Environ Health Perspect*. 2004;112(8):879–82.

17. Chan JK, Fanucchi MV, Anderson DS, Abid AD, Wallis CD, Dickinson DA, Kumfer BM, Kennedy IM, Wexler AS, Van Winkle LS. Susceptibility to inhaled flame-generated ultrafine soot in neonatal and adult rat lungs. *Toxicol Sci.* 2011;124(2):472–86.
18. Choi JH, Won MH. Microglia in the normally aged hippocampus. *Lab Anim Res.* 2011;27(3):181–7.
19. Church JS, Chace-Donahue F, Blum JL, Ratner JR, Zelikoff JT, Schwartzter JJ. Neuroinflammatory and behavioral outcomes measured in adult offspring of mice exposed prenatally to e-cigarette aerosols. *Environ Health Perspect.* 2020;128(4): 047006.
20. Church JS, Tamayo JM, Ashwood P, Schwartzter JJ. Repeated allergic asthma in early versus late pregnancy differentially impacts offspring brain and behavior development. *Brain Behav Immun.* 2021;93:66–79.
21. Church JS, Tijerina PB, Emerson FJ, Coburn MA, Blum JL, Zelikoff JT, Schwartzter JJ. Perinatal exposure to concentrated ambient particulates results in autism-like behavioral deficits in adult mice. *Neurotoxicology.* 2018;65:231–40.
22. Claeson AS, Palmquist E, Lind N, Nordin S. Symptom-trigger factors other than allergens in asthma and allergy. *Int J Environ Health Res.* 2016;26(4):448–57.
23. Clougherty JE, Levy JJ, Kubzansky LD, Ryan PB, Suglia SF, Canner MJ, Wright RJ. Synergistic effects of traffic-related air pollution and exposure to violence on urban asthma etiology. *Environ Health Perspect.* 2007;115(8):1140–6.
24. Conti E, Mitra J, Calderoni S, Pannek K, Shen KK, Pagnozzi A, Rose S, Mazzotti S, Scelfo D, Tosetti M, et al. Network over-connectivity differentiates autism spectrum disorder from other developmental disorders in toddlers: a diffusion MRI study. *Hum Brain Mapp.* 2017;38:2333–44.
25. Croen LA, Grether JK, Yoshida CK, Odouli R, Van de Water J. Maternal autoimmune diseases, asthma and allergies, and childhood autism spectrum disorders: a case-control study. *Arch Pediatr Adolesc Med.* 2005;159(2):151–7.
26. Croen LA, Qian Y, Ashwood P, Zerbo O, Schendel D, Pinto-Martin J, Daniele Fallin M, Levy S, Schieve LA, Yeargin-Allsopp M, Sabourin KR, Ames JL. Infection and fever in pregnancy and autism spectrum disorders: findings from the study to explore early development. *Autism Res.* 2019;12(10):1551–61.
27. Damianidou E, Mouratidou L, Kyrousi C. Research models of neurodevelopmental disorders: The right model in the right place. *Front Neurosci.* 2022;18:46.
28. Davoli-Ferreira M, Thomson CA, McCoy KD. Microbiota and microglia interactions in ASD. *Front Immunol.* 2021;12: 676255.
29. De Araujo EG, Da Silva GM, Dos Santos AA. Neuronal cell survival: the role of interleukins. *Ann NY Acad Sci.* 2009;1153(1):57–64.
30. Diz-Chaves Y, Astiz M, Bellini MJ, Garcia-Segura LM. Prenatal stress increases the expression of proinflammatory cytokines and exacerbates the inflammatory response to LPS in the hippocampal formation of adult male mice. *Brain Behav Immun.* 2013;28:196–206.
31. Ecker C. The neuroanatomy of autism spectrum disorder: an overview of structural neuroimaging findings and their translatability to the clinical setting. *Autism.* 2017;21(1):18–28.
32. Eggen BJL, Raj D, Hanisch UK, Boddeke HWGM. Microglial phenotype and adaptation. *J Neuroimmune Pharmacol.* 2013;8:807–23.
33. Fasmer OB, Halmøy A, Eagan TM, Oedegaard KJ, Haavik J. Adult attention deficit hyperactivity disorder is associated with asthma. *BMC Psychiatry.* 2011;11:128.
34. Fazel N, Kundi M, Jensen-Jarolim E, Pali-Schöll I, Kazemzadeh A, Abdizadeh MF, Esmaily H, Akbarzadeh R, Ahmadi R. Prospective cohort study of pregnancy complications and birth outcomes in women with asthma. *Arch Gynecol Obstet.* 2018;298(2):279–87.
35. Filiano AJ, Xu Y, Tustison NJ, Marsh RL, Baker W, Smirnov I, Overall CC, Gadani SP, Turner SD, Weng Z, Peerzade SN, Chen H, Lee KS, Scott MM, Beenhakker MP, Litvak V, Kipnis J. Unexpected role of interferon- γ in regulating neuronal connectivity and social behaviour. *Nature.* 2016;535(7612):425–9.
36. Flood L, Korol SV, Ekselius L, Birnir B, Jin Z. Interferon- γ potentiates GABAA receptor-mediated inhibitory currents in rat hippocampal CA1 pyramidal neurons. *J Neuroimmunol.* 2019;337: 577050.
37. Garay PA, Hsiao EY, Patterson PH, McAllister AK. Maternal immune activation causes age- and region-specific changes in brain cytokines in offspring throughout development. *Brain Behav Immun.* 2013;31:54–68.
38. Giovanoli S, Weber-Stadlbauer U, Schedlowski M, Meyer U, Engler H. Prenatal immune activation causes hippocampal synaptic deficits in the absence of overt microglia anomalies. *Brain Behav Immun.* 2016;55:25–38.
39. Gluck JC, Gluck PA. The effect of pregnancy on the course of asthma. *Immunol Allergy Clin N Am.* 2006;26(1):63–80.
40. Goines PE, Croen LA, Braunschweig D, Yoshida CK, Grether J, Hansen R, Kharrazi M, Ashwood P, Van de Water J. Increased midgestational IFN- γ , IL-4 and IL-5 in women bearing a child with autism: a case-control study. *Mol Autism.* 2011;2:13.
41. Gong C, Chu M, Yang J, Gong X, Han B, Chen L, Bai Z, Wang J, Zhang Y. Ambient fine particulate matter exposures and human early placental inflammation. *Environ Pollut.* 2022;315: 120446.
42. Gong T, Lundholm C, Rejnö G, Bölte S, Larsson H, D'Onofrio BM, Lichtenstein P, Almqvist C. Parental asthma and risk of autism spectrum disorder in offspring: a population and family-based case-control study. *Clin Exp Allergy.* 2019;49(6):883–91.
43. Groen W, Teluij M, Buitelaar J, Tendolkar I. Amygdala and hippocampus enlargement during adolescence in autism. *J Am Acad Child Adolesc Psychiatry.* 2010;49(6):552–60.
44. Guarneri M, Balmes JR. Outdoor air pollution and asthma. *Lancet (London, England).* 2014;383(9928):1581–92.
45. Guglielmotti C, Le Blon D, Santermans E, Salas-Perdomo A, Daans J, De Vocht N, Shah D, Hoornaert C, Praet J, Peerlings J, Kara F, Bigot C, Mai Z, Goossens H, Hens N, Hendrix S, Verhoye M, Planas AM, Berneman Z, van der Linden A, et al. Interleukin-13 immune gene therapy prevents CNS inflammation and demyelination via alternative activation of microglia and macrophages. *Glia.* 2016;64(12):2181–200.
46. Han VX, Patel S, Jones HF, Nielsen TC, Mohammad SS, Hofer MJ, Gold W, Brilot F, Lain SJ, Nassar N, Dale RC. Maternal acute and chronic inflammation in pregnancy is associated with common neurodevelopmental disorders: a systematic review. *Transl Psychiatry.* 2021;11(1):71.
47. Hisle-Gorman E, Susi A, Stokes T, Gorman G, Erdie-Lalena C, Nylund CM. Prenatal, perinatal, and neonatal risk factors of autism spectrum disorder. *Pediatr Res.* 2018;84(2):190–8.
48. Hopkins LE, Laing EA, Peake JL, Uyeminami D, Mack SM, Li X, Smiley-Jewell S, Pinkerton KE. Repeated iron-soot exposure and nose-to-brain transport of inhaled ultrafine particles. *Toxicol Pathol.* 2018;46(1):75–84.
49. Hua L, Ju L, Xu H, Li C, Sun S, Zhang Q, Cao J, Ding R. Outdoor air pollution exposure and the risk of asthma and wheezing in the offspring. *Environ Sci Pollut Res Int.* 2023;30(6):14165–89.
50. Hughes HK, Moreno RJ, Ashwood P. Innate immune dysfunction and neuroinflammation in autism spectrum disorder (ASD). *Brain Behav Immun.* 2023;108:245–54.
51. Hughes HK, Ashwood P. Overlapping evidence of innate immune dysfunction in psychotic and affective disorders. *Brain, behavior, & immunity - health.* 2020;2: 100038.
52. Hughes HK, Mills Ko E, Rose D, Ashwood P. Immune dysfunction and autoimmunity as pathological mechanisms in autism spectrum disorders. *Front Cell Neurosci.* 2018;12:405.
53. Jacquet A. Innate immune responses in house dust mite allergy. *ISRN Allergy.* 2013;2013: 735031.
54. Jasinski J, Pinkerton KE, Kennedy IM, Leppert VJ. Spatially resolved energy electron loss spectroscopy studies of iron oxide nanoparticles. *Microsc Microanal.* 2006;12(5):424–31.
55. Jiang HY, Xu LL, Shao L, Xia RM, Yu ZH, Ling ZX, Yang F, Deng M, Ruan B. Maternal infection during pregnancy and risk of autism spectrum disorders: a systematic review and meta-analysis. *Brain Behav Immun.* 2016;58:165–72.
56. Juckel G, Manitz MP, Brüne M, Friebe A, Heneka MT, Wolf RJ. Microglial activation in a neuroinflammatory animal model of schizophrenia—a pilot study. *Schizophr Res.* 2011;131(1–3):96–100.
57. Kalkbrenner AE, Windham GC, Serre ML, Akita Y, Wang X, Hoffman K, Thayer BP, Daniels JL. Particulate matter exposure, prenatal and postnatal windows of susceptibility, and autism spectrum disorders. *Epidemiology.* 2015;26(1):30–42.
58. Kebir H, Kreyenborg K, Ifergan I, Dodelet-Devillers A, Cayrol R, Bernard M, Giuliani F, Arbour N, Becher B, Prat A. Human TH17 lymphocytes

- promote blood–brain barrier disruption and central nervous system inflammation. *Nat Med.* 2007;13(10):1173–5.
59. Knuesel I, Chicha L, Britschgi M, Schobel SA, Bodmer M, Hellings JA, et al. Maternal immune activation and abnormal brain development across CNS disorders. *Nat Rev Neurol.* 2014;10:643–60.
 60. Koyama R, Ikegaya Y. Microglia in the pathogenesis of autism spectrum disorders. *Neurosci Res.* 2015;100:1–5.
 61. Lazic SE, Essioux L. Improving basic and translational science by accounting for litter-to-litter variation in animal models. *BMC Neurosci.* 2013;14:37.
 62. Lazic SE, Clarke-Williams CJ, Munafò MR. What exactly is “N” in cell culture and animal experiments? *PLoS Biol.* 2018;16(4): e2005282.
 63. Le Blon D, Guglielmetti C, Hoornaert C, Quarta A, Daans J, Dooley D, Lemmens E, Praet J, De Vocht N, Reekmans K, Santermans E, Hens N, Goossens H, Verhoye M, Van der Linden A, Berneman Z, Hendrix S, Ponsaerts P. Intracerebral transplantation of interleukin 13-producing mesenchymal stem cells limits microgliosis, oligodendrocyte loss and demyelination in the cuprizone mouse model. *J Neuroinflamm.* 2016;13(1):288.
 64. Leaderer BP, Belanger K, Triche E, Holford T, Gold DR, Kim Y, Jankun T, Ren P, McSharry JE, Platts-Mills TA, Chapman MD, Bracken MB. Dust mite, cockroach, cat, and dog allergen concentrations in homes of asthmatic children in the northeastern United States: impact of socioeconomic factors and population density. *Environ Health Perspect.* 2002;110(4):419–25.
 65. Lee J, Kim SJ, Son TG, Chan SL, Mattson MP. Interferon-gamma is up-regulated in the hippocampus in response to intermittent fasting and protects hippocampal neurons against excitotoxicity. *J Neurosci Res.* 2006;83(8):1552–7.
 66. Lock C, Hermans G, Pedotti R, Brendolan A, Schadt E, Garren H, Langer-Gould A, Strober S, Cannella B, Allard J, Klonowski P, Austin A, Lad N, Kaminski N, Galli SJ, Oksenberg JR, Raine CS, Heller R, Steinman L. Gene-microarray analysis of multiple sclerosis lesions yields new targets validated in autoimmune encephalomyelitis. *Nat Med.* 2002;8(5):500–8.
 67. Longhin E, Holme JA, Gualtieri M, Camatini M, Øvrevik J. Milan winter fine particulate matter (wPM2.5) induces IL-6 and IL-8 synthesis in human bronchial BEAS-2B cells, but specifically impairs IL-8 release. *Toxicol In Vitro.* 2018;52:365–73.
 68. Lyall K, Ashwood P, Van de Water J, Hertz-Picciotto I. Maternal immune-mediated conditions, autism spectrum disorders, and developmental delay. *J Autism Dev Disord.* 2014;44(7):1546–55.
 69. Ma L, Li XW, Zhang SJ, Yang F, Zhu GM, Yuan XB, Jiang W. Interleukin-1 beta guides the migration of cortical neurons. *J Neuroinflamm.* 2014;11:114.
 70. Macia L, Viltart O, Delacore M, Sachot C, Héliot L, Di Santo JP, Wolowczuk I. Interleukin-7, a new cytokine targeting the mouse hypothalamic arcuate nucleus: role in body weight and food intake regulation. *PLoS ONE.* 2010;5(4): e9953.
 71. Martel MJ, Rey E, Beauchesne MF, Malo JL, Perreault S, Forget A, Blais L. Control and severity of asthma during pregnancy are associated with asthma incidence in offspring: two-stage case-control study. *Eur Respir J.* 2009;34(3):579–87.
 72. Matsui EC, Hansel NN, McCormack MC, Rusher R, Breyse PN, Diette GB. Asthma in the inner city and the indoor environment. *Immunol Allergy Clin N Am.* 2008;28(3):665–x.
 73. Maximo J, Cadena EJ, Kana RK. The implications of brain connectivity in the neuropsychology of autism. *Neuropsychol Rev.* 2014;24:16–31.
 74. Mayer A, Ulrich A, Czerwinski J, Mooney J. Metal-Oxide Particles in Combustion Engine Exhaust. SAE Technical Paper 2010-01-0792, 2010.
 75. Miao W, Zhao Y, Huang Y, Chen D, Luo C, Su W, Gao Y. IL-13 ameliorates neuroinflammation and promotes functional recovery after traumatic brain injury. *J Immunol (Baltimore, Md.: 1950).* 2020;204(6):1486–98.
 76. Michaelson MD, Mehler MF, Xu H, Gross RE, Kessler JA. Interleukin-7 is trophic for embryonic neurons and is expressed in developing brain. *Dev Biol.* 1996;179(1):251–63.
 77. Mills BD, Pearce HL, Khan O, Jarrett BR, Fair DA, Lahvis GP. Prenatal domoic acid exposure disrupts mouse pro-social behavior and functional connectivity MRI. *Behav Brain Res.* 2016;308:14–23.
 78. Milovanovic J, Arsenijevic A, Stojanovic B, Kanjevac T, Arsenijevic D, Radosavljevic G, Milovanovic M, Arsenijevic N. Interleukin-17 in chronic inflammatory neurological diseases. *Front Immunol.* 2020;11:947.
 79. Mitschik S, Schierl R, Nowak D, Jörres RA. Effects of particulate matter on cytokine production in vitro: a comparative analysis of published studies. *Inhalation Toxicol.* 2008;20(4):399–414.
 80. Modabbernia A, Velthorst E, Reichenberg A. Environmental risk factors for autism: an evidence-based review of systematic reviews and meta-analyses. *Mol Autism.* 2017;8:13.
 81. Morato Torres CA, Wassouf Z, Zafar F, Sastre D, Outeiro TF, Schüle B. The role of alpha-synuclein and other Parkinson’s genes in neurodevelopmental and neurodegenerative disorders. *Int J Mol Sci.* 2020;21(16):5724.
 82. Morgan JT, Barger N, Amaral DG, Schumann CM. Stereological study of amygdala glial populations in adolescents and adults with autism spectrum disorder. *PLoS ONE.* 2014;9(10): e110356.
 83. Morgan JT, Chana G, Pardo CA, Achim C, Semedferi K, Buckwalter J, et al. Microglial activation and increased microglial density observed in the dorsolateral prefrontal cortex in autism. *Biol Psychiatry.* 2010;68(4):368–76.
 84. Mortimer K, Neugebauer R, Lurmann F, Alcorn S, Balmes J, Tager I. Air pollution and pulmonary function in asthmatic children: effects of prenatal and lifetime exposures. *Epidemiology.* 2008;19(4):550–62.
 85. Murphy VE. Asthma in pregnancy—management, maternal co-morbidities, and long-term health. *Best practice and research. Clin Obstet Gynaecol.* 2022;85(Pt A):45–56.
 86. Murphy CM, Deeley Q, Daly EM, Ecker C, O’Brien FM, Hallahan B, Loth E, Toal F, Reed S, Hales S, Robertson DM, Craig MC, Mullins D, Barker GJ, Lavender T, Johnston P, Murphy KC, Murphy DG. Anatomy and aging of the amygdala and hippocampus in autism spectrum disorder: an in vivo magnetic resonance imaging study of Asperger syndrome. *Autism Res.* 2012;5(1):3–12.
 87. Murphy VE, Clifton VL, Gibson PG. Asthma exacerbations during pregnancy: incidence and association with adverse pregnancy outcomes. *Thorax.* 2006;61(2):169–76.
 88. Murphy VE, Gibson P, Talbot PI, Clifton VL. Severe asthma exacerbations during pregnancy. *Obstet Gynecol.* 2005;106(5 Pt 1):1046–54.
 89. Musah S, DeJarnett N, Hoyle GW. Tumor necrosis factor- α mediates interactions between macrophages and epithelial cells underlying proinflammatory gene expression induced by particulate matter. *Toxicology.* 2012;299(2–3):125–32.
 90. Ottum PA, Arellano G, Reyes LI, Iruretagoyena M, Naves R. Opposing roles of interferon-gamma on cells of the central nervous system in autoimmune neuroinflammation. *Front Immunol.* 2015;6:539.
 91. Pan HC, Yang CN, Hung YW, Lee WJ, Tien HR, Shen CC, Sheehan J, Chou CT, Sheu ML. Reciprocal modulation of C/EBP- α and C/EBP- β by IL-13 in activated microglia prevents neuronal death. *Eur J Immunol.* 2013;43(11):2854–65.
 92. Park SY, Kang MJ, Han JS. Interleukin-1 beta promotes neuronal differentiation through the Wnt5a/RhoA/JNK pathway in cortical neural precursor cells. *Mol Brain.* 2018;11(1):39.
 93. Patel S, Dale RC, Rose D, Heath B, Nordahl CW, Rogers S, Guastella AJ, Ashwood P. Maternal immune conditions are increased in males with autism spectrum disorders and are associated with behavioural and emotional but not cognitive co-morbidity. *Transl Psychiatry.* 2020;10(1):286.
 94. Patterson PH. Immune involvement in schizophrenia and autism: etiology, pathology and animal models. *Behav Brain Res.* 2009;204(2):313–21.
 95. Pinkerton KE, Zhou Y, Zhong C, Smith KR, Teague SV, Kennedy IM, Ménache MG. Mechanisms of particulate matter toxicity in neonatal and young adult rat lungs. *Res Rep (Health Effects Institute).* 2008;135:3–52.
 96. Popa M, Peltecu G, Gica N, Ciobanu AM, Botezatu R, Gica C, Steriade A, Panaitescu AM. Asthma in pregnancy. Review of current literature and recommendations. *Maedica.* 2021;16(1):80–7.
 97. Pui DYH, Chen SC, Zuo Z. PM2.5 in China: measurements, sources, visibility and health effects, and mitigation. *Particuology.* 2014;13:1–26.
 98. Rasband WS. ImageJ. U. S. National Institutes of Health, Bethesda, Maryland, USA.
 99. Raz R, Roberts AL, Lyall K, Hart JE, Just AC, Laden F, Weisskopf MG. Autism spectrum disorder and particulate matter air pollution before, during, and after pregnancy: a nested case-control analysis

- within the Nurses' Health Study II Cohort. *Environ Health Perspect.* 2015;123(3):264–70.
100. Renz H, Holt PG, Inouye M, Logan AC, Prescott SL, Sly PD. An exposome perspective: Early-life events and immune development in a changing world. *J Allerg Clin Immunol* 2017;140(1):24–40.
 101. Réus GZ, Fries GR, Stertz L, Badawy M, Passos IC, Barichello T, et al. The role of inflammation and microglial activation in the pathophysiology of psychiatric disorders. *Neuroscience.* 2015;300:141–54.
 102. Richards R, Greimel E, Kliemann D, Koerte IK, Schulte-Körne G, Reuter M, Wachinger C. Increased hippocampal shape asymmetry and volumetric ventricular asymmetry in autism spectrum disorder. *NeuroImage Clin.* 2020;26: 102207.
 103. Roberts AL, Lyall K, Hart JE, Laden F, Just AC, Bobb JF, Koenen KC, Ascherio A, Weisskopf MG. Perinatal air pollutant exposures and autism spectrum disorder in the children of Nurses' Health Study II participants. *Environ Health Perspect.* 2013;121(8):978–84.
 104. Rojas DC, Peterson E, Winterrowd E, Reite ML, Rogers SJ, Tregellas JR. Regional gray matter volumetric changes in autism associated with social and repetitive behavior symptoms. *BMC Psychiatry.* 2006;6:56.
 105. Rossignol DA, Genuis SJ, Frye RE. Environmental toxicants and autism spectrum disorders: a systematic review. *Transl Psychiatry.* 2014;4(2): e360.
 106. Salo PM, Arbes SJ Jr, Crockett PW, Thorne PS, Cohn RD, Zeldin DC. Exposure to multiple indoor allergens in US homes and its relationship to asthma. *J Allergy Clin Immunol.* 2008;121(3):678–684.e2.
 107. Schielzeth H, Nakagawa S. Nested by design: model fitting and interpretation in a mixed model era. *Methods Ecol Evol.* 2013;4:14–24.
 108. Schraufnagel DE. The health effects of ultrafine particles. *Exp Mol Med.* 2020;52(3):311–7.
 109. Schwartzter JJ, Careaga M, Chang C, Onore CE, Ashwood P. Allergic fetal priming leads to developmental, behavioral and neurobiological changes in mice. *Transl Psychiatry.* 2015;5(4): e543.
 110. Schwartzter JJ, Careaga M, Coburn MA, Rose DR, Hughes HK, Ashwood P. Behavioral impact of maternal allergic-asthma in two genetically distinct mouse strains. *Brain Behav Immun.* 2017;63:99–107.
 111. Silbajoris R, Osornio-Vargas AR, Simmons SO, Reed W, Bromberg PA, Dailey LA, Samet JM. Ambient particulate matter induces interleukin-8 expression through an alternative NF- κ B (nuclear factor-kappa B) mechanism in human airway epithelial cells. *Environ Health Perspect.* 2011;119(10):1379–83.
 112. Ślusarczyk J, Trojan E, Głombik K, Budziszewska B, Kubera M, Lasoń W, Popiołek-Barczyk K, Mika J, Wędzony K, Basta-Kaim A. Prenatal stress is a vulnerability factor for altered morphology and biological activity of microglia cells. *Front Cell Neurosci.* 2015;9:82.
 113. Steiner S, Bisig C, Petri-Fink A, Rothen-Rutishauser B. Diesel exhaust: current knowledge of adverse effects and underlying cellular mechanisms. *Arch Toxicol.* 2016;90(7):1541–53.
 114. Straughen JK, Sitarik AR, Johnson CC, Wegienka G, Ownby DR, Johnson-Hooper TM, Allo G, Levin AM, Cassidy-Bushrow AE. Prenatal IgE as a risk factor for the development of childhood neurodevelopmental disorders. *Front Pediatr.* 2021;9: 601092.
 115. Tamayo JM, Rose D, Church JS, Schwartzter JJ, Ashwood P. Maternal allergic asthma induces prenatal neuroinflammation. *Brain Sci.* 2022;12(8):1041.
 116. Tioleco N, Silberman AE, Stratigos K, Banerjee-Basu S, Spann MN, Whitaker AH, Turner JB. Prenatal maternal infection and risk for autism in offspring: a meta-analysis. *Autism Res.* 2021;14(6):1296–316.
 117. Vogel Ciernia A, Careaga M, LaSalle JM, Ashwood P. Microglia from offspring of dams with allergic asthma exhibit epigenomic alterations in genes dysregulated in autism. *Glia.* 2018;66(3):505–21.
 118. Volk HE, Hertz-Picciotto I, Delwiche L, Lurmann F, McConnell R. Residential proximity to freeways and autism in the CHARGE study. *Environ Health Perspect.* 2011;119(6):873–7.
 119. Volk HE, Lurmann F, Penfold B, Hertz-Picciotto I, McConnell R. Traffic-related air pollution, particulate matter, and autism. *JAMA Psychiat.* 2013;70(1):71–7.
 120. Volk HE, Park B, Hollingue C, Jones KL, Ashwood P, Windham GC, Lurman F, Alexeef SE, Kharrazi M, Pearl M, Van de Water J, Croen LA. Maternal immune response and air pollution exposure during pregnancy: insights from the Early Markers for Autism (EMA) study. *J Neurodev Disord.* 2020;12(1):42.
 121. von Ehrenstein OS, Aralis H, Cockburn M, Ritz B. In utero exposure to toxic air pollutants and risk of childhood autism. *Epidemiology.* 2014;25(6):851–8.
 122. Wainwright PE, Leatherdale ST, Dubin JA. Advantages of mixed effects models over traditional ANOVA models in developmental studies: a worked example in a mouse model of fetal alcohol syndrome. *Dev Psychobiol.* 2007;49(7):664–74.
 123. Wang P, You D, Saravia J, Shen H, Cormier SA. Maternal exposure to combustion generated PM inhibits pulmonary Th1 maturation and concomitantly enhances postnatal asthma development in offspring. *Part Fibre Toxicol.* 2013;10:29.
 124. Yagishita K, Suzuki R, Mizuno S, Katoh-Semba R, Sadakata T, Sano Y, et al. CAPS2 deficiency affects environmental enrichment-induced adult neurogenesis and differentiation/survival of newborn neurons in the hippocampal dentate gyrus. *Neurosci Lett.* 2017;661:121–5.
 125. Zablotzky B, Black LI, Maenner MJ, Schieve LA, Danielson ML, Bitsko RH, Blumberg SJ, Kogan MD, Boyle CA (2019). Prevalence and trends of developmental disabilities among children in the United States: 2009–2017. *Pediatrics* 2019;144(4).
 126. Zhong CY, Zhou YM, Smith KR, Kennedy IM, Chen CY, Aust AE, Pinkerton KE. Oxidative injury in the lungs of neonatal rats following short-term exposure to ultrafine iron and soot particles. *J Toxicol Environ Health A.* 2010;73(12):837–47.
 127. Zhou YM, Zhong CY, Kennedy IM, Leppert VJ, Pinkerton KE. Oxidative stress and NF κ B activation in the lungs of rats: a synergistic interaction between soot and iron particles. *Toxicol Appl Pharmacol.* 2003;190(2):157–69.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

